



# LYMPHA PRESS ORDER FORM

DATE OF ORDER \_\_\_\_\_

Medical Solutions Supplier is the exclusive provider of Lympha Press.

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_ Phone \_\_\_\_\_

### Extremities for Treatment

LEGS LT RT With Abdomen (Lympha Pants)

ARMS LT RT With Trunk (Comfy Sleeve)

Special Request \_\_\_\_\_

Treatment Pressures \_\_\_\_\_ Frequency \_\_\_\_\_ / Day \_\_\_\_\_ MIN Length of Necessity \_\_\_\_\_ Months (99=lifetime)

Default Settings 40mmHg, TID/BID, 60 min

### Diagnosis

- Q82.0 HEREDITARY LYMPHEDEMA
- I89.0 SECONDARY LYMPHEDEMA DUE TO:
  - VENOUS INSUFFICIENCY
  - TUMOR(S)
  - SCARRING OF THE LYMPH CHANNELS FROM CELLULITIS AND/OR LYMPHANGITIS
  - CANCER
- I97.2 POST MASTECTOMY SYNDROME
- I87.2 CHRONIC VENOUS INSUFFICIENCY
  - VARICOSE VEIN WITH ULCER
  - VENOUS HYPERTENSION WITH ULCER

### Prescribing Physician

Last \_\_\_\_\_ First \_\_\_\_\_

NPI# \_\_\_\_\_

Phone \_\_\_\_\_ Fax \_\_\_\_\_

Email \_\_\_\_\_

Physician Signature \_\_\_\_\_

In Clinic Contact \_\_\_\_\_

Date \_\_\_\_\_

# Medicare Requires Two Office Visits 4-5 Weeks Apart.

## First Office Visit Documentation:

### Diagnosis of Lymphedema

- > Hereditary (Q82.0)
- > Secondary (I89.0)
- > Post mastectomy (I97.2)

### Limb Circumference Measurements

- > Ankle and calf for lower extremity lymphedema
- > Wrist and forearm for upper extremity lymphedema

## Start (or continuation of) all of the following conservative care treatments:

- > Class 2 stockings/sleeves or compression wraps
- > Elevation of the affected limb(s)
- > Exercise (ankle pumps, walking, etc.)

## Second Office Visit Documentation (4-5 weeks later):

- > Continued diagnosis of lymphedema
- > Limb circumference measurements
- > Patient compliance with compression system, elevation, and exercise
- > Significant lymphedema symptoms persist, or conservative care did not significantly improve symptoms

## Lympha Press Contraindications

Use of the Lympha Press® system is not recommended in the presence of one or more of the following conditions:

- Known or suspected deep vein thrombosis (DVT) or pulmonary embolism
- During the inflammatory phlebitis process
- Acute infection of the affected limb
- Decompensated cardiac failure or other ischemic vascular disease
- Severe arteriosclerosis or other ischemic vascular disease
- Any circumstance where increased venous and lymphatic return is undesirable
- The abdominal area should not be treated during pregnancy

## Sample Dictations

Patient presents today with stage 2 lymphedema. Will begin (or continue) conservative treatments including elevation, exercise, and class 2 compression stockings. *Include circumferential limb measurements.*

### After 4 weeks (28-42 days)

Patient presents today with stage 2 lymphedema. Despite compliance with conservative treatments including elevation, exercise, and class 2 compression stockings, the patient has not shown significant improvement, and symptoms of chronic and severe lymphedema persist. *Include circumferential limb measurements.*



lymphapress.com  
Questions? Call 800-734-0422